

VCH-PHC PHARMACY DEPARTMENT ANTIBIOTIC CROSS-ALLERGY CHART

	Amikacin	Amoxicillin	Ampicillin	Azithromycin	Cefadroxil (NF)	Cefazolin	Cefepime	Cefixime	Cefotaxime	Cefoxitin	Cefprozil (NF)	Ceftazidime	Ceftriaxone	Cefuroxime	Cephalexin	Chloramphenicol	Ciprofloxacin	Clarithromycin	Clindamycin	Cloxacillin	Cotrimoxazole (Sulfa)	Daptomycin	Ertapenem	Erythromycin	Gentamicin	Imipenem	Levofloxacin	Meropenem	Metronidazole	Moxifloxacin	Penicillin	Piperacillin/Tazobactam	Streptomycin	Tigecycline	Tobramycin	Vancomycin			
Amikacin																																							
Amoxicillin																																							
Ampicillin		X																																					
Azithromycin																																							
Cefadroxil (NF)		X	X																																				
Cefazolin																																							
Cefepime																																							
Cefixime																																							
Cefotaxime																																							
Cefoxitin																																							
Cefprozil (NF)		X	X		X																																		
Ceftazidime																																							
Ceftriaxone																																							
Cefuroxime																																							
Cephalexin		X	X		X																																		
Chloramphenicol																																							
Ciprofloxacin																																							
Clarithromycin																																							
Clindamycin																																							
Cloxacillin																																							
Cotrimoxazole (Sulfa)																																							
Daptomycin																																							
Ertapenem																																							
Erythromycin																																							
Gentamicin		X																																					
Imipenem																																							
Levofloxacin																																							
Meropenem																																							
Metronidazole																																							
Moxifloxacin																																							
Penicillin		X	X																																				
Piperacillin/Tazobactam		X	X		*																																		
Streptomycin		X																																					
Tigecycline																																							
Tobramycin		X																																					
Vancomycin																																							

KEY TO SYMBOLS:

Blank = Not cross-sensitive as different structure

X = **AVOID.** Potential for cross-sensitivity due to similar structure or based on clinical reports of cross-sensitivity

* = **USE WITH CAUTION.** Similar side chain - Potential for cross reaction. Monitor first dose closely (vital signs at 15, 30, and 60 minutes post-first dose)

NF = Non-formulary drug

Note: **AVOID all penicillins, cephalosporins and carbapenems** if patient reports history of **severe delayed skin allergy to ANY beta-lactam antibiotic** (i.e. Steven-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug rash with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP)).